

Glucose Monitoring System Quick Operating Guide for Blood Sample



1 The Home screen



2 Wash patient's hand thoroughly and massage finger to stimulate blood flow.



3 Insert a test strip into the meter. A test strip with a blood drop will display.



4 Use a safety lancet to puncture the finger.



5 Squeeze the finger to form a drop of blood. Wipe away the first drop of blood then squeeze the finger again to form a second drop of blood.



6 When the blood drop appears, touch the end of the test strip to the blood drop until the test strip fills and the meter beeps.

7 Warning!

The test strip must fill completely upon touching the blood droplet. If the test strip does not fill completely, do not touch the blood droplet a second time. Discard the test strip and repeat the test with a new test strip.



8 Glucose test results are available on-screen in 6 seconds. There is one long beep when the results are ready. There are 3 short beeps if test results are outside the range of the test strip.

- 9 Remove the test strip and discard into an appropriate biohazard container.
- 10 When patient testing is complete, the StatStrip Xpress2 Glucose Hospital Meter should be cleaned and disinfected after use prior to testing with a new patient. For instructions, see Cleaning and Disinfecting the Meter section in the Instructions for Use Manual.
- 11 Test results are automatically saved. If no activity for 1 minute, the meter will time-out: screen goes blank.



Nova Biomedical, 200 Prospect Street, Waltham, MA 02454 Tel: 800-545-6682 • www.novabiomedical.com